
Fludarabine Actavis 50

fludarabine phosphate 50 mg powder for injection vial

Consumer Medicine Information

May 2009

What is in this leaflet?

This leaflet answers some common questions about Fludarabine Actavis 50.

It does not contain all of the available information.

It does not take the place of talking to your doctor.

All medicines have risks and benefits. Your doctor has weighed the risks of using Fludarabine Actavis 50 against the benefits it is expected to have for you. Only your doctor is able to weigh up all the relevant facts and you should consult him/her about all aspects of this medication as it relates to you.

If you have any concerns about using Fludarabine Actavis 50, ask your doctor.

Keep this leaflet. You may want to read it again.

What Fludarabine Actavis 50 is used for and how it works

Fludarabine Actavis 50 is used to treat a form of leukaemia known as chronic lymphocytic leukaemia, or CLL.

All cells in the body produce new cells like themselves by dividing. For this purpose, the cells' genetic material (DNA) must be copied and reproduced. Fludarabine Actavis 50 works by hindering the production of new DNA. Therefore, when Fludarabine Actavis 50 is taken up by the cancer cells it stops the growth of new cancer cells. It has been discovered that Fludarabine Actavis 50 works especially well against some cancers of the white blood cells.

Before you are given Fludarabine Actavis 50

When you must not use it

You must not be given Fludarabine Actavis 50 if any of the following apply to you:

- Pregnancy or breast-feeding
- Allergy (hypersensitivity) to any of the ingredients of this medicine
- Your kidney function is severely reduced
- The number of red blood cells is reduced due to a breaking down of these cells (hemolytic anemia)

The effects of Fludarabine Actavis 50 have not been studied in children. It should not be used together with the leukaemia drug pentostatin.

Fludarabine Actavis 50

fludarabine phosphate 50 mg Powder for injection(50mg/vial)

Consumer Information Leaflet

Before you use it

If it is found that your kidneys do not work properly you may be given this medicine at a reduced dose.

If your liver does not work properly, you have a poor state of health or if you are over 75 years old, Fludarabine Actavis 50 will be given with caution.

Men, and women who may still be fertile, must use a reliable form of contraception during and for at least 6 months after stopping treatment

Check with your doctor about any vaccinations you may need, because live vaccinations should be avoided during and after treatment with Fludarabine Actavis 50.

It is not known whether Fludarabine Actavis 50 decreases your fertility.

Fludarabine Actavis 50 may reduce your ability to drive or use machines, since tiredness, weakness and visual disturbances have been observed. Patients experiencing such adverse effects should avoid driving and using machines.

While using Fludarabine Actavis 50

When used in patients with acute leukemia at doses four times greater than the recommended dose for chronic lymphocytic leukemia (CLL), a third of patients experienced severe central nervous system effects including blindness, coma and death. Such effects have only rarely (coma and agitation) or uncommonly (confusion) been reported in patients who receive the recommended dose for CLL. However, you should mention to your doctor any unusual

symptoms. The effect of long-term use of Fludarabine Actavis 50 on the central nervous system is unknown.

The number of normal blood cells may also be reduced for periods of up to 12 months duration. You will have regular blood tests during treatment. On some occasions death has been reported in patients experiencing reductions in the number of normal blood cells.

Special regular checkups will be necessary in case you are over 75 years old, or your liver or kidneys do not work properly.

If you need a blood transfusion and you are being (or have been) treated with Fludarabine Actavis 50, you should mention this to the doctor.

Fludarabine Actavis 50

fludarabine phosphate 50 mg Powder for injection(50mg/vial)

Consumer Information Leaflet

If your disease is very severe, your body may not be able to get rid of all the waste products from the cells destroyed by Fludarabine Actavis 50. This may cause dehydration, kidney failure and heart problems. Your doctor will be aware of this and may give you other drugs to stop this happening.

The worsening or flare up of pre-existing skin cancer as well as new onset of skin cancer has been reported in patients during or after Fludarabine Actavis 50 therapy.

Taking other medicines

This medicine should not be used with another drug called pentostatin (deoxycoformycin).

The effectiveness of Fludarabine Actavis 50 may be reduced by medications containing dipyridamole and similar substances. If you are

taking any other medicines regularly, including medicines bought from a supermarket, health food store or pharmacy, tell your doctor.

How Fludarabine Actavis 50 is given

How much is given

The dose of Fludarabine Actavis 50 is calculated according to your body surface, which is calculated from your weight and height.

The recommended dose is 25 mg fludarabine phosphate/m² body surface given daily for 5 consecutive days every 28 days by the intravenous route.

How it is given

Fludarabine Actavis 50 is given as a 5 day treatment course which is repeated each month. The treatment involves

injecting Fludarabine Actavis 50 into a vein (often in the arm) once each day for 5 consecutive days. Your specialist, who should have experience with similar medications, will work out what dose is right for you.

How long it is given

Your doctor will decide how long to continue your treatment with Fludarabine Actavis 50.

Unwanted Effects

Like many leukaemia drugs Fludarabine Actavis 50 is a very strong drug with many possible side effects, many of which can be very serious. Fludarabine Actavis 50 can harm normal blood cells as well as the CLL blood cells, resulting in severe anaemia, abnormal bleeding and bruising and serious infections, especially pneumonia.

Fludarabine Actavis 50

fludarabine phosphate 50 mg Powder for injection(50mg/vial)

Consumer Information Leaflet

Fludarabine Actavis 50 can also commonly cause:

- fever and chills,
- infection,
- generally feeling unwell,
- nausea and vomiting,
- tiredness,
- weakness,
- numb or weak limbs,
- build up of fluid in the body tissues,
- visual problems,
- loss of appetite,
- inflammation of the lining of the mouth (stomatitis) and gut,
- skin rashes.

An allergic reaction in the lungs, symptoms of which are difficulties in breathing and coughing can also rarely occur. If you experience any difficulty in breathing or have a cough tell your doctor immediately.

If you notice any unusual bruising, excessive bleeding after injury or if you develop signs of

infections (e.g fever, chills, feeling unwell, pain), motor or visual disturbances, tell your doctor immediately.

Further side effects that have occurred in patients with chronic lymphatic leukaemia treated with Fludarabine Actavis 50 are:

- peripheral neuropathy (disturbances of the nervous system of your body);
- diarrhoea,
- mucositis
- stomatitis and uncommonly gastrointestinal bleeding;
- rare cases of haemorrhagic cystitis (inflammation with associated bleeding from the mucosa of the bladder);
- tumour lysis syndrome.

If you notice a pain in your side, blood in your

urine or that your stools are tar-coloured or covered with blood tell your doctor immediately.

In rare cases heart failure, irregular heart beat and skin cancer have been reported in patients treated with Fludarabine Actavis 50.

Do not be alarmed by this list of possible side effects. You may not experience any of them. This is not a complete list of all possible side effects.

You should immediately report any side effects that you notice to your specialist or to the nursing staff.

Overdosage

The exact dose of Fludarabine Actavis 50 is worked out by your specialist. The dose is determined by your weight and height.

Fludarabine Actavis 50

fludarabine phosphate 50 mg Powder for injection(50mg/vial)

Consumer Information Leaflet

Overdosage is unlikely to occur. In case of an overdose your doctor will stop the therapy and treat the symptoms.

After using Fludarabine Actavis 50

Storage

Fludarabine Actavis 50 will usually be stored at the hospital below 25°C and protected from light.

- Fludarabine Actavis 50 should be stored where children cannot reach it or see it.
- Fludarabine Actavis 50 must not be used after the expiry date stated on the box and on the vial.

Disposal

Any unused medicine must be disposed appropriately by the medical staff.

Product Description

Fludarabine Actavis 50 is a white or almost white powder in a glass vial. It is available as 50 mg/vial.

Each vial is for a single injection only. Fludarabine Actavis 50 has to be reconstituted before use.

Ingredients

Fludarabine Actavis 50 contains 50 mg Fludarabine phosphate. Other ingredients in the vial are mannitol and sodium hydroxide.

Sponsor

Actavis Australia Pty Ltd
63 Old Street
North Adelaide
South Australia 5006
ABN 43 122 896 468
Telephone Number:
1300 881 893

Distributor in Australia:

Solely distributed in Australia by:
Generic Health Pty Ltd.,

Suite 1, 1175 Toorak Road,
Camberwell VIC 3124.

Australian Registration Number:

Fludarabine Actavis 50 fludarabine phosphate 50 mg powder for injection vial:
AUST R 147831

This leaflet was prepared in May 2009.