

CEFTRIAXONE INJECTION

(Max Pharma Pty Ltd)

CEFTRIAXONE SODIUM

CONSUMER MEDICINE INFORMATION

WHAT IS IN THIS LEAFLET

This leaflet answers some common questions about CEFTRIAXONE INJECTION. It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist. All medicines have risks and benefits. Your doctor has weighed the risks of you being given CEFTRIAXONE INJECTION against the benefits they expect it will have for you. If you have any concerns about being given this medicine, ask your doctor or pharmacist.

Keep this leaflet. You may need to read it again.

WHAT CEFTRIAXONE INJECTION IS USED FOR

CEFTRIAXONE INJECTION is an antibiotic used to treat infections in different parts of the body caused by bacteria.

CEFTRIAXONE INJECTION is also used to prevent infections before, during and after surgery.

CEFTRIAXONE INJECTION will not work against infections caused by viruses such as colds or the flu.

CEFTRIAXONE INJECTION belongs to a group of antibiotics called cephalosporins. These antibiotics work by killing the bacteria that are causing your infection or by stopping bacteria from growing.

Your doctor may have prescribed CEFTRIAXONE INJECTION for another reason.

Ask your doctor if you have any questions why CEFTRIAXONE INJECTION has been prescribed for you.

This medicine is available only with a doctor's prescription.

CEFTRIAXONE INJECTION is not addictive.

BEFORE YOU ARE GIVEN CEFTRIAZONE INJECTION

When you must not be given it

Do not use CEFTRIAZONE INJECTION if:

- You have had an allergic reaction to ceftriazone, or any other cephalosporins.

Some of the symptoms of an allergic reaction may include severe skin rash, itching, hives, dry skin, swelling of the face, lips, mouth or throat which may cause difficulty in swallowing or breathing, swelling of the hands, feet or ankles.

- You have had a serious allergic reaction to penicillins. You may have an increased chance of being allergic to CEFTRIAZONE INJECTION if you are allergic to penicillins.

In newborns, CEFTRIAZONE INJECTION and IV calcium-containing solutions must not be given within 48 hours of each other.

If you are not sure if you should start using CEFTRIAZONE INJECTION, contact your doctor.

Before you are given it

Your doctor must know about all the following before you are given CEFTRIAZONE INJECTION.

Tell your doctor if you have allergies to:

1. Penicillin medicines. You may have an increased chance of being allergic to CEFTRIAZONE INJECTION if you are allergic to penicillins.
2. Lignocaine or other anaesthetic drugs. These substances may be used to dissolve CEFTRIAZONE INJECTION before administration.
3. Any other medicines.
4. Any other substances including foods, preservatives or dyes.

Tell your doctor if you are pregnant, intend to become pregnant, are breast feeding or plan to breastfeed. Your doctor will discuss the risks and benefits of using CEFTRIAZONE INJECTION during pregnancy or when breastfeeding.

If you have not told your doctor about any of the above, tell them before you are given CEFTRIAZONE INJECTION.

Tell your doctor if you have or have ever had any medical conditions, especially the following:

- kidney disease
- bowel conditions e.g. inflammation of the large bowel.

TAKING OTHER MEDICINES

Tell your doctor if you are taking any other medicines including any that you buy without a prescription from your pharmacy, supermarket or health food shop.

In particular tell your doctor if you are using chloramphenicol eye drops (Chloromycetin, Chlorsig). This medicine may be affected by CEFTRIAZONE INJECTION, or may affect how well it works. You may need to take different amounts of your medicine, or you may need to take different medicines. Your doctor will advise you.

Talk to your doctor about the need for an additional method of contraception while using CEFTRIAZONE INJECTION.

Some antibiotics may decrease the effectiveness of some birth control pills, although this has not been shown with CEFTRIAZONE INJECTION.

Your doctor and pharmacist may have more information on medicines to be careful with or avoid while being given CEFTRIAZONE INJECTION.

HOW CEFTRIAZONE INJECTION IS GIVEN

CEFTRIAZONE INJECTION is given by a slow injection into a vein. CEFTRIAZONE INJECTION must only be given by a doctor or nurse.

CEFTRIAZONE INJECTION should not be mixed or administered with calcium-containing solutions, even via different infusion lines.

Your doctor will decide what dose and how long you will receive CEFTRIAZONE INJECTION. This depends on your infection and other factors, such as your weight. For most infections, CEFTRIAZONE INJECTION is usually given once a day. The length of treatment usually is 4-14 days.

Sometimes only a single dose of CEFTRIAZONE INJECTION is required for the treatment and prevention of certain infections.

If too much CEFTRIAZONE INJECTION is given, you may experience symptoms such as skin reactions, chills, diarrhoea, stomach upsets, headache or dizziness.

WHILE YOU ARE USING CEFTRIAZONE INJECTION

Things you must do

- If the symptoms of your infection do not improve within a few days, or if they become worse, tell your doctor.
- If you get severe diarrhoea tell your doctor, pharmacist or nurse immediately. Do this even if it occurs several weeks after CEFTRIAZONE INJECTION has been stopped. Diarrhoea may mean that you have a serious condition affecting your bowel. You may need urgent medical care. Do not take any diarrhoea medicine without first checking with your doctor.
- If you get a sore or furry white mouth or tongue while using or soon after stopping CEFTRIAZONE INJECTION, tell your doctor. Also tell your doctor if you get vaginal itching or discharge. This may mean that you have a fungal infection called thrush. Sometimes the use of CEFTRIAZONE INJECTION allows fungi to grow and the above symptoms to occur. CEFTRIAZONE INJECTION does not work against fungi.
- If you become pregnant while using CEFTRIAZONE INJECTION, tell your doctor.
- If you have to test your urine for sugar while you are being given CEFTRIAZONE INJECTION, make sure your doctor knows which type of test you use. CEFTRIAZONE INJECTION may affect the results of some of these tests.
- If you have any blood tests tell your doctor you are being given CEFTRIAZONE INJECTION. CEFTRIAZONE INJECTION may affect the results of some blood tests.

Tell all doctors, dentists and pharmacists who are treating you that you are using CEFTRIAZONE INJECTION.

Things you must not do

- Do not give CEFTRIAZONE INJECTION to anyone else even if they seem to have the same condition as you.
- Do not use CEFTRIAZONE INJECTION to treat other complaints unless your doctor says to.

Things to be careful of

Be careful driving or operating machinery until you know how CEFTRIAZONE INJECTION affects you. CEFTRIAZONE INJECTION generally does not cause any problems with your ability to drive or operate machinery. However, as with many other medicines, CEFTRIAZONE INJECTION may cause dizziness in some people.

Side Effects

Tell your doctor or pharmacist as soon as possible if you do not feel well while you are using CEFTRIAXONE INJECTION.

CEFTRIAXONE INJECTION helps most people with infections but it may have unwanted side effects in some people. All medicines can have unwanted effects. Sometimes they are serious, most of the time they are not. Do not be alarmed by this list of possible side effects. You may not experience any of them.

You may need medical treatment if you get some of the side effects. **Ask your doctor or pharmacist to answer any questions you may have.**

While using it

Tell your doctor if you notice any of the following and they worry you:

- white, furry, sore tongue and mouth (oral thrush)
- sore and itchy vagina and/or discharge (vaginal thrush)
- a hard lump, swelling, pain or tenderness at the injection site
- diarrhoea, feeling sick, vomiting
- altered sense of taste
- headache, dizziness
- fast or irregular heart beat.

Tell your doctor immediately, or go to Accident and Emergency at your nearest hospital if you notice any of the following:

- severe skin rash, itching or hives
- red, peeling or dry skin
- swelling of the face, lips, mouth or throat which may cause difficulty in swallowing or breathing, swelling of the hands, feet or ankles.

These may be serious side effects of CEFTRIAXONE INJECTION. You may need urgent medical attention. Serious side effects are rare.

After finishing it

Tell your doctor immediately if you notice any of the following side effects, particularly if they occur several weeks after stopping treatment with CEFTRIAXONE INJECTION:

- severe abdominal cramps or stomach cramps
- watery and severe diarrhoea
- raised temperature (fever) in combination with one or both of the above.

These are rare but serious side effects. You may have a serious condition affecting your bowel. Therefore, you may need urgent medical attention. However, this side effect is rare.

Do not take any diarrhoea medicine without first checking with your doctor.

Ask your doctor or pharmacist if you don't understand anything in this list.

Other side effects not listed above may also occur in some patients. Tell your doctor if you notice any other side effects.

PRODUCT DESCRIPTION

Storage

CEFTRIAXONE INJECTION will be stored in the pharmacy or on the ward. The powder for injection is kept in a cool dry place where the temperature stays below 25°C.

What CEFTRIAXONE INJECTION looks like

CEFTRIAXONE INJECTION is available as a white to yellowish-orange powder. It is packaged in a clear glass vial. It is reconstituted with a suitable fluid before injection into a vein.

Ingredients

CEFTRIAXONE INJECTION contains the active ingredient, ceftriaxone as the sodium salt.

- each 1 gram CEFTRIAXONE INJECTION vial contains 1 gram ceftriaxone

CEFTRIAXONE INJECTION 1 gram is available in packs of 5 vials.

SPONSOR

Lupin Australia
Suite 5, 310-314 Whitehorse Road
Balwyn Victoria 3103
Australia

DISTRIBUTOR

Max Pharma Pty Ltd
Suite 10, 400 Canterbury Road
Surrey Hills Victoria 3127
Australia

This leaflet was prepared in December 2008

Australian Registration Numbers:

CEFTRIAXONE INJECTION
1 gram vial: AUST R 104489

A40001/2